

Literatur

- 1 **Gonthier A, Favrat B. Syndrome de fatigue chronique. Rev Med Suisse. 2015 Nov;496:2236-42.
- 2 *Davis HE, McCorkell L, Vogel JM, et al. Long COVID: major findings, mechanisms and recommendations. Nat Rev Microbiol. 2023 Mar;21(3):133-46.
- 3 Choutka J, Jansari V, Hornig M, et al. Unexplained post-acute infection syndromes. Nat Med. 2022 May;28(5):911-23.
- 4 **National Institute for Health and Care Excellence guideline. Myalgic encephalomyelitis (or encephalopathy)/chronic fatigue syndrome: diagnosis and management. London: NICE; 2021.
- 5 Appelman B, Charlton BT, Goulding RP, et al. Muscle abnormalities worsen after post-exertional malaise in long COVID. Nat Commun. 2024 Jan 4;15:17.
- 6 Gupta G, Buonsenso D, Wood J, et al. Mechanistic Insights Into Long Covid: Viral Persistence, Immune Dysregulation, and Multi-Organ Dysfunction. Compr Physiol. 2025 Jun;15(3):e70019.
- 7 Vernon SD, Hartle M, Sullivan K, et al. Post-exertional malaise among people with long COVID compared to myalgic encephalomyelitis/chronic fatigue syndrome (ME/CFS). Work. 2023;74(4):1179-86.
- 8 **Stussman B, Williams A, Snow J, et al. Characterization of Post-exertional Malaise in Patients With Myalgic Encephalomyelitis/Chronic Fatigue Syndrome. Front Neurol. 2020 Sep 18;11:1025.
- 9 *Davis HE, Assaf GS, McCorkell L, et al. Characterizing long COVID in an international cohort: 7 months of symptoms and their impact. EClinicalMedicine. 2021 Aug;38:101019.
- 10 Soini E, Rosenström T, Määttänen I, et al. Physical activity and specific symptoms of depression: A pooled analysis of six cohort studies. J Affect Disord. 2024 Mar 1;348:44-53.
- 11 *Vallée G, Xi D, Avramovic G, et al. Evaluating the longitudinal physical and psychological health effects of persistent long Covid 3.5 years after infection. PLOS One. 2025 Jun 24;20(6):e0326790.
- 12 König RS, Paris DH, Sollberger M, et al. Identifying the mental health burden in Myalgic Encephalomyelitis/Chronic Fatigue Syndrome (ME/CFS) patients in Switzerland: A pilot study. Heliyon. 2024 Feb 23;10(5):e27031.
- 13 Boulazreg S, Rokach A. The Lonely, Isolating, and Alienating Implications of Myalgic Encephalomyelitis/Chronic Fatigue Syndrome. Healthcare (Basel). 2020 Oct 20;8(4):413.
- 14 Gómez Bravo R, Infanti A, Billieux J, et al. Unmasking the Psychological Landscape of Long COVID: A Cluster-analytical Approach. Biopsychosoc Sci Med. 2025 Apr 1;87(3):162-72.
- 15 Taylor RR, Jason LA, Curie CJ. Prognosis of Chronic Fatigue in a Community-Based Sample. Psychosom Med. 2002 Mar-Apr;64(2):319-27.
- 16 Guggisberg J, Liechti L, Keller T, et al. Auswirkungen von Long-Covid auf die Invalidenversicherung [Online]. 1. Januar 2025. Verfügbar unter: www.buerobass.ch/fileadmin/Files/2025/BSV_2025_Covid-Monitoring_Forschungsbericht_d.pdf
- 17 Grach SL, Seltzer J, Chon TY, et al. Diagnosis and Management of Myalgic Encephalomyelitis/Chronic Fatigue Syndrome. Mayo Clin Proc. 2023 Oct;98(10):1544-51.
- 18 *Goudsmit EM, Nijs J, Jason LA, et al. Pacing as a strategy to improve energy management in myalgic encephalomyelitis/chronic fatigue syndrome: a consensus document. Disabil Rehabil. 2012;34(13):1140-7.
- 19 Cairns R, Hotopf M. A systematic review describing the prognosis of chronic fatigue syndrome. Occup Med (Lond). 2005 Jan;55(1):20-31.
- 20 *Ballouz T, Menges D, Anagnostopoulos A, et al. Recovery and symptom trajectories up to two years after SARS-CoV-2 infection: population based, longitudinal cohort study. BMJ. 2023 May 31;381:e074425.

- 21 *Bundesamt für Sozialversicherungen. Versicherte mit Long-Covid in der Invalidenversicherung (IV) [Online]. (abgerufen am 31. Januar 2026). Verfügbar unter:
www.bsv.admin.ch/de/versicherte-long-covid-iv
- 22 Chu L, Valencia IJ, Garvert DW, et al. Deconstructing post-exertional malaise in myalgic encephalomyelitis/chronic fatigue syndrome: A patient-centered, cross-sectional survey. PLoS ONE. 2018 Jun 1;13(6):e0197811.
- 23 Holtzman CS, Bhatia S, Cotler J, et al. Assessment of Post-Exertional Malaise (PEM) in Patients with Myalgic Encephalomyelitis (ME) and Chronic Fatigue Syndrome (CFS): A Patient-Driven Survey. Diagnostics (Basel). 2019 Mar 2;9(1):26.
- 24 Batheja S, Yang J, Liao Y, et al. Fatigue in major depression is linked to depression severity, not sleep duration or anterior cingulate cortex neurochemistry. J Affect Disord. 2026 Apr 15;399:121076.
- 25 Szuhany KL, Simon NM. Anxiety Disorders: A Review. JAMA. 2022 Dec 27;328(24):2431-45.
- 26 Ghanean H, Ceniti AK, Kennedy SH. Fatigue in Patients with Major Depressive Disorder: Prevalence, Burden and Pharmacological Approaches to Management. CNS Drugs. 2018 Jan;32(1):65-74.
- 27 Cotler J, Holtzman C, Dudun C, et al. A Brief Questionnaire to Assess Post-Exertional Malaise. Diagnostics (Basel). 2018 Sep 11;8(3):66.
- 28 Nugent K, Berdine G. Dyspnea and long COVID patients. Am J Med Sci. 2024 Oct;368(4):399-404.
- 29 Mancini DM, Brunjes DL, Cook D, et al. Abnormal breathing patterns and hyperventilation are common in patients with chronic fatigue syndrome during exercise. Front Med (Lausanne). 2025 Nov 10;12:1669036.

* lesenswert

** unbedingt lesenswert